

9700207

THE UNITED STAYES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Pioneer Hi-Bred International, Inc.

MICROS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR PORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT AD BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PH02T'

In Destinant Thereof, I have hereunto set my hand and caused the seal of the Hunt Inviety Hvotection Office to be affixed at the City of Washington, D.C. this fifth day of Tebruary, in the year of our Lord two thousand one.

alanktont

Arting Commissioner Plant Variety Protection Office Agricultural Marketing Service Sary of Agriculture

CD 470 mt as: 15

14A. Exhibit A. Origin and Breeding History

Pedigree: PHR03/PHWT8)X31K13K3X

Pioneer Line PH02T, Zea mays L., a yellow corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross PHR03 X PHWT8 using the pedigree method of breeding. The progenitors of PH02T are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing and selection were practiced within the above F1 cross for 6 generations in the development of PH02T at Windfall, Indiana. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Windfall, Indiana, as well as other Pioneer research stations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations made for uniformity.

PH02T has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity.

No variant traits have been observed or are expected in PH02T.

The criteria used in the selection of PH02T were yield, both per se and in hybrid combinations; kernel size, especially important in production; ability to germinate in adverse conditions; number of tillers, especially important in production because having numerous tillers increases hybrid production costs spent on detasseling; disease and insect resistance; pollen yield; tassel size and pollen shed duration.

JUS 100

Variety PHWT8 was derived by pedigree selection from PHZ51 (PVP Certificate Number 8600132) X PHW89. Variety PHW89 was derived by pedigree selection from PHG44 X G35 (PVP Certificate Number 8300140). Variety PHG44 was derived by pedigree selection from PH891 X 207 (PVP Certificate Number 8300144). Variety PH891 was derived by pedigree selection. The majority of the pedigree relative contribution to the derivation of PH891 was B14, A556, A509, I205, IDT, I205, MINN49, Troyer Reid.

DEVELOPMENTAL HISTORY FOR PH02T

Season/Year	Inbreeding Level
Summer 1990	FO
Winter 1990	F1
Summer 1991	F2*
Summer 1992	F3*
Winter 1992	F4*
Summer 1993	F5*
Winter 1993	F6*
Summer 1994	F7*
Winter 1994	F8**

^{*}PH02T was selfed and selected through F7 generation.

 $[\]ensuremath{^{\star\star}\text{PH02T}}$ was selfed and ear-rowed for F8 generation.

Exhibit B Novelty Statement

The data in Table 1A and 1B are from paired comparisons collected primarily in Johnston, IA in 1997. The traits in Table 1A, 1B and Exhibit D collectively show measurable differences between the two varieties.

PH02T mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PHBG4 (PVP Certificate No. 9500202).

Variety PH02T has shorter husk extension length (2.5 cm vs 5.7 cm) than PHBG4 (Table 1A, 1B).

Variety PH02T has shorter husk length (23.7 cm vs 26.4 cm) than PHBG4 (Table 1A, 1B).

Variety PH02T has longer kernel length (11.0 mm vs 9.9 mm) than PHBG4 (Table 1A, 1B).

Variety PH02T has more total leaves (19.8 leaves per plant vs 18.2 leaves per plant) than PHBG4 (Table 1A, 1B).

Variety PH02T has a shorter tassel central spike length (20.7 cm vs 29.7 cm) than PHBG4 (Table 1A, 1B).

Variety PH02T has a shorter tassel length (48.8 cm vs 58.8 cm) than PHBG4 (Table 1A, 1B).

Variety PH02T has more primary tassel branches (11.2 vs 6.9) than PHBG4 (Table 1A, 1B).

Variety PH02T reaches 50% pollen shed (GDUSHD) later than PHBG4 (1573 GDU's vs 1542 GDU's) (Exhibit D).

Variety PH02T reaches 50% silking (GDUSLK) later than PHBG4 (1590 GDU's vs 1557 GDU's) (Exhibit D).

Exhibit B Novelty Statement Tables

Table 1A. Data from two environments in Johnston, IA in 1997 are supporting evidence for differences between PH02T and PHBG4

<u>8</u>	Staff loc year on	vanety- 1	Variety≟ variety- C 1 2	Count- Count- 1 2	ount-M	eant-1	Weah-1 Mean-2 Mean Diff	Mean Diff	DF Pooled	l Value Pooled	Prob (2- tail) Pooled
	1997 husk extension length (cm)	PH02T	PHBG4	2	5	2.2	5.6	-3.4	8	-6.21	0.000
	1997 husk extension length (cm)	PH02T	PHBG4	ည	5	2.8	5.8	-3.0	æ	-4.87	0.001
	1997 husk length (cm)	PH02T	PHBG4	5	5	24.4	26.8	-2.4	\times	-5.37	0.001
	1997 husk length (cm)	PH02T	PHBG4	5	2	23.0	26.0	-3.0	∞	-5.48	0.001
	1997 kernel length (mm)	PH02T	PHBG4	5	5	11.2	10.0	1.2	æ	2.45	0.040
	1997 kernel length (mm)	PH02T	PHBG4	5	2	10.8	9.8	1.0	œ	2.36	0.046
	1997 leaf number (# of leaves/plant)	PH02T	PHBG4	5	5	20.0	18.4	1.6	∞	4.00	
	1997 leaf number (# of leaves/plant)	PH02T	PHBG4	2	ည	19.6	18.0	1.6	æ	2.67	0.029
	1997 tassel central spike length (cm)	PH02T	PHBG4	5	5	22.6	29.6	-7.0	Φ	4.50	0.002
	1997 tassel central spike length (cm)	PH02T	PHBG4	2	2	18.8	29.8	-11.0	8	-6.60	0.000
	1997 tassel length (cm)	PH02T	PHBG4	5	2	52.8	59.6	9.9	x	-2.27	0.053
	1997 tassel length (cm)	PH02T	PHBG4	က	က	44.8	58.0	-13.2	∞	-5.01	0.001
	1997 tassel primary branch (# of primary	PH02T	PHBG4	5	2	11.6	6.8	4.8	ω	4.80	0.001
	branches)		10.0.00m.	***************************************							:
	1997 tassel primary branch (# of primary PH02T	PH02T	PHBG4	5	ည	10.8	7.0	3.8	∞	3.92	0.004
	branches)								and the state of t		A COLUMN TO SERVICE STORY

Table 1B. Summary data across environments in 1997 are supporting evidence for differences between PH02T and PHBG4. Locations had different environmental conditions.

Prob_(2- tail)_Poole d	0.000	0.000	0.003	0.000	0.000	0.000	0.000	
d Fro	95	31	50	54	-7.21	31	41	
t-Value_ Pooled_	-7.	φ	က်	4.	-7.	4	Ö	
DF_ Pooled	18	18	18	18	18	18	18	
vean- Mean_Diff	-3.2	-2.7	=	1.6	-9.0	-10.0	4.3	mayor Auri Mila
Mean⊦ 2	5.7	26.4	9.9	18.2	29.7	58.8	6.9	
Mean-1	2.5	23.7	11.0	19.8	20.7	48.8	11.2	
Sunf-2	5	10	10	10	10	10	10	A
variety-2 Count-1 Count-2	10	10	10	10	10	10	10	
variety-2	PHBG4	PHBG4	PHBG4	PHBG4	PHBG4	PHBG4	PHBG4	
variety-1	PH02T	PH02T	PH02T	PH02T	PH02T	PH02T	PH02T	
ar Fait	1997 husk extension length (cm)	1997 husk length (cm)	1997 kernel length (mm)	1997 leaf number (# of leaves/plant)	1997 tassel central spike length (cm)	1997 tassel length (cm)	1997 tassel primary branch (# of	primary branches)
ye								

Exhibit C (Corn: Maize)

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

Objective Description of Variety Corn (Zea mays L.)

Name of Applicant (s)		Variety Seed Source	Variety Name or To	emporary Designation
Pioneer Hi-Bred	International, Inc.		PH027	Г
Address (Street & No.,	or RFD No., City, State, Zip Cod	le and Country	FOR OFFICIAL USE	31
7301 NW 62 nd Av	venue, P.O. Box 85,		9	700207
Johnston, Iowa	•		PVP0 Number	700207
Place the appropriate n	umber that describes the varietal of	characters typical of this inbred	variety in the spaces be	low. Right justify whole
	ling zeroes if necessary. Complete			
	considered necessary for an ade			
	se in conjunction with Munsell co			d #26 in Comments section):
01=Light Green	06=Pale Yellow	11=Pink	16=Pale Purple	21=Buff
02=Medium Green	07=Yellow	12=Light Red	17=Purple	22=Tan
03=Dark Green	08=Yellow Orange	13=Cherry Red	18=Colorless	23=Brown
04=Very Dark Green	09=Salmon	14=Red	19=White	24=Bronze
05=Green-Yellow	10=Pink-Orange	15=Red & White	20=White Capped	25=Variegated (Describe)
				26=Other (Describe)
STANDARD INBRED	CHOICES			
(Use the most similar (i	n background and maturity) of th	ese to make comparisons based	d on grow-out trial data	a):
Yellow Dent Families:		Yellow Dent (Unrelated): Sweet C	Corn:
Family Members		Co109, ND246,	C13, I	owa5125, P39, 2132
B14 CM105, A	632, B64, B68	Oh7, T232,		
B37, B76,	H84	W117, W153R,	Popcorr	a:
B73 N192, A67	9, B73, NC268	W18BN	SG153	33, 4722, HP301, HP7211
C103 Mo17, Val	02, Va35, A682			
Oh43 A619, MS7	71, H99, Va26	White Dent:	Pipecor	n:
WF9 W64A, A5	54, A654, Pa91	C166, H105, Ky228	Mo15	W, Mo16W, Mo24W

Ceres/worddata/doug/96pvp

TYPE: (describe intermediate types in Comments section):			Standa	rd Inbred	Name
2 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental			<u> </u>	<u>'A26</u>	
2. REGION WHERE DEVELOPED IN THE U.S.A.:		· ··	Standa	rd Seed S	Source
4 1=Northwest 2=Northcentral 3=Northeast 4=Southeast 5=Southeast 5	uthcentral		ļ ,	MES 193	329
6=Southwest 7=Other			_	WILO 150	<u> </u>
3. MATURITY (In Region of Best Adaptability; show Heat Unit formula in	'Comments' se	ection)			
DAYS HEAT UNITS			DAYS F	HEAT UN	ITS
076 1,537.5 From emergence to 50% of plants in silk			069	<u>1,352.5</u>	
077 1,561.3 From emergence to 50% of plants in pollen			070	1,378.3	
005 0,119.8 From 10% to 90% pollen shed			007	0,162.5	
From 50% silk to optimum edible quality					
067 1,163.5 From 50% silk to harvest at 25% moisture			068	<u>1,279.8</u>	
4. PLANT:	Standard	Sample	5	Standard	Samp
	Deviation	Size	r	Deviation	Size
221.3 cm Plant Height (to tassel tip)	08.22	<u>04</u>	<u>215.5</u>	11.82	<u>04</u>
084.0 cm Ear Height (to base of top ear node)	04.32	04	076.5	11.70	04
016.4 cm Length of Top Ear Internode	01.17	04	013.3	01.14	04
Ø _∞ ∂ -1:0 Average Number of Tillers	00.00	04	0.01.0	<u>08.68</u>	04
1.0 Average Number of Ears per Stalk	00.00	04	1.0	00.00	04
3 Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate	e 4=Dark		1		
5. LEAF:	Standard	Sample		Standard	Sampl
	Deviation	Size	Ĺ	Deviation	Size
10.4 cm Width of Ear Node Leaf	01.00	<u>04</u>	08.9	00.19	<u>04</u>
82.2 cm Length of Ear Node Leaf	02.08	<u>04</u>	69.7	<u>07.18</u>	<u>04</u>
06 Number of leaves above top ear	00.34	04	<u>06</u>	00.26	<u>04</u>
30 Degrees Leaf Angle (measure from 2nd leaf above ear at anthesis to stalk above leaf)	<u>12.99</u>	<u>04</u>	<u>35</u>	12.40	<u>04</u>
03 Leaf Color (Munsell code) 5GY34			03	<u>5G\</u>	<u> </u>
1 Leaf Sheath Pubescence (Rate on scale from 1=none to 9=lik	e peach fuzz)		1		
6 Marginal Waves (Rate on scale from 1=none to 9=many)			<u>8</u>		
6 Longitudinal Creases (Rate on scale from 1=none to 9=many)			<u>8</u> <u>8</u>		
6. TASSEL:	Standard	Sample		Standard	Samp
	Deviation	Size	[Deviation	Size
09 Number of Primary Lateral Branches	02.36	<u>04</u>	<u>18</u>	01.43	<u>04</u>
47 Branch Angle from Central Spike	<u>13.28</u>	<u>04</u>	<u>63</u>	<u> 19.92</u>	<u>04</u>
52.5 cm Tassel Length (from top leaf collar to tassel tip)	02.22	<u>04</u>	<u>61.2</u>	<u>02.89</u>	<u>04</u>
8 Pollen Shed (rate on scale from 0=male sterile to 9=heavy she	ed)		<u>8</u>		
11 Anther Color (Munsell code) 7.5R68			<u>07</u>	<u>10`</u>	<u> 488</u>
01 Glume Color (Munsell code) 5GY66			01	<u>5G</u>	<u>Y58</u>
1 Bar Glumes (Glume Bands): 1=Absent 2=Present			1		
			L		

5/5/00

Application	Nariety Data	PH02T	Page 2			Standar	d Inbred	Data
7a. EAR	(Unhusked Data):					01		
<u>01</u>	Silk Color (3 days at	fter emergence) (Mu	unsell code)		2.5GY86	707	2.5G	<u> 186</u>
<u>02</u>	Fresh Husk Color (2	5 days after 50% sil	king) (Munsell code)		5GY68	<u>02</u>	5GY	<u>56</u>
<u>21</u>	Dry Husk Color (65	days after 50% silkir	ng) (Munseli code)		2.5Y84	<u>21</u>	2.5Y8	3.52
2	Position of Ear at Dr	y Husk Stage: 1= U	pright 2= Horizontal	3= Pendant		<u>3</u>		
<u>5</u>	Husk Tightness (Rat	te of Scale from 1=v	ery loose to 9=very t	ight)		<u>5</u>		
<u>2</u>	Husk Extension (at I	narvest): 1=Short (ea	ars exposed) 2=Med	ium (<8 cm)		3		
	3=Long (8-10 cm be	yond ear tip) 4=Very	/ Long (>10 cm)					
7b. EAR	(Husked Ear Data):			Standard	Sample	Stan	dard	Sample
				Deviation	Size	Devi	ation	Size
<u>17.1</u>	cm Ear Length			00.41	<u>04</u>	<u>13.4</u> 0	1.08	<u>04</u>
<u>41.4</u>	mm Ear Diameter at	t mid-point		02.60	<u>04</u>	<u>42.3</u> 0	0.53	<u>04</u>
<u>132.2</u>	gm Ear Weight			20.24	<u>04</u>	<u>104.3</u> 0	7.69	<u>04</u>
<u>15</u>	Number of Kernel R	ows		00.86	<u>04</u>	<u>15.8</u> 0	0.60	<u>04</u>
<u>2</u>	Kernel Rows: 1=Ind	istinct 2=Distinct				<u>2</u>		
<u>1</u>	Row Alignment: 1=5	Straight 2=Slightly C	urved 3=Spiral			1 1		
<u>10.7</u>	cm Shank Length			01.47	<u>04</u>	<u>09.8</u> 0	0.85	<u>04</u>
<u>2</u>	Ear Taper: 1=Slight	2= Average 3=Extre	eme			2		
8. KERNI	EL (Dried)			Standard	Sample	Standa	rd	Sample
				Deviation	Size	Deviati	on	Size
10.9	mm Kernel Length			00.98	<u>04</u>	10.8 <u>0</u>	0.30	<u>04</u>
<u>07.6</u>	mm Kernel Width			00.49	<u>04</u>	08.1 0	0.12	<u>04</u>
04.8	mm Kernel Thickness	5		00.28	<u>04</u>	04.0 0	<u>0.10</u>	<u>04</u>
<u>19.6</u>	% Round Kernels (Si	nape Grade)		06.59	<u>04</u>	16.0 O	5.44	<u>04</u>
1	Aleurone Color Patte	rn: 1-Homozygous:	2=Segregating			1		
<u>07</u>	Aluerone Color (Mur	sell code)		<u>10</u>	YR712	<u>07</u>	2.5Y8	<u>314</u>
<u>07</u>	Hard Endosperm Col	or (Munsell code)		<u>10`</u>	YR714	<u>07</u>	2.5Y8	<u>314</u>
<u>03</u>	Endosperm Type:					<u>3</u>		
	4=High Amylose S	=Extra Sweet (sh2) Starch 5=Waxy Star Super Sweet (se)	ch 6=High Protein					
<u>27.0</u>	gm Weight per 100 K	 ernels (unsized sam	nple)	04.69	<u>04</u>	<u>24.25</u> 0	2.06	<u>04</u>
9. COB:		·		Standard	Sample	Sta	andard	Sampl
				Deviation	Size	-	viation	Size
<u>23.</u> 4	mm Cob Diameter at	mid-point		01.87	<u>04</u>	<u>25.5</u> (01.25	<u>04</u>
	Cob Color (Munsell c	•	<u>10R58</u>			<u>19</u>	2.5	

Application Variety Data

Page 3

Standard Inbred Data

	RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant); I	eave blank
if not tested	l; leave Race or Strain Options blank if polygenic):	
A. Leaf E	Blights, Wilts, and Local Infection Diseases	
	Anthracnose Leaf Blight (Colletotrichum graminicola)	
7	Common Rust (Puccinia sorghi)	<u>6</u>
-	Common Smut (Ustilago maydis)	_
	Eyespot (Kabatiella zeae)	
	Goss's Wilt (Clavibacter michiganense spp. nebraskense)	
<u>6</u>	Gray Leaf Spot (Cercospora zeae-maydis)	2
~	Helminthosporium Leaf Spot (Bipolaris zeicola) Race ——	_
<u>3</u>	Northern Leaf Blight (Exserohilum turcicum) Race ———	3
<u>5</u>	Southern Leaf Blight (Bipolaris maydis) Race ——	3
-	Southern Rust (Puccinia polysora)	_
<u>8</u>	Stewart's Wilt (Erwinia stewartii)	4
-	Other (Specify) ———	_
B. Syster	nic Diseases	
<u>5</u>	Corn Lethal Necrosis (MCMV and MDMV)	<u>3</u>
4	Head Smut (Sphacelotheca reiliana)	6
	Maize Chlorotic Dwarf Virus (MDV)	
	Maize Chlorotic Mottle Virus (MCMV)	
<u>4</u>	Maize Dwarf Mosaic Virus (MDMV)	<u>3</u>
_	Sorghum Downy Mildew of Corn (Peronosclerospora sorghi)	_
	Other (Specify) ———	
C. Stalk i	Rots	
. <u>5</u>	Anthracnose Stalk Rot (Colletotrichum graminicola)	<u>3</u>
2	Diplodia Stalk Rot (Stenocarpella maydis)	2
	Fusarium Stalk Rot (Fusarium moniliforme)	
	Gibberella Stalk Rot (Gibberella zeae)	
	Other (Specify)	
D. Ear an	d Kernel Rots	
	Aspergillus Ear and Kernel Rot (Aspergillus flavus)	
<u>3</u>	Diplodia Ear Rot (Stenocarpella maydis)	<u>5</u>
<u>3</u>	Fusarium Ear and Kernel Rot (Fusarium moniliforme)	<u>5</u>
_	Gibberella Ear Rot (Gibberella zeae)	
	Other (Specify) ———	

Application Variety Data

Page 3

Standard Inbred Data

PH02T

Application Variety Data

Page 4

Standard Inbred Dat

CLARIFICATION OF DATA IN EXHIBITS C AND D

Please note the data presented in Exhibit C, "Objective Description of Variety," is data collected primarily at Johnston, Iowa plus description information from the maintaining station. The data in Exhibit D, "Additional Description of Variety," is data from comparisons of inbreds grown in the same tests in the adapted growing area of PH02T.

5MS 7/27/00 The data collected in exhibit C was collected in 1995 and 1996 for page 1 and 2. There are environmental factors that differ from year to year. In 1995, May was wet and August was warmer. In 1996, May was very wet and August was cool with very little heat or drought stress compared to most years. Environmental temperature and precipitation differences during the vegetative and grain fill periods can impact plant and grain traits and be a source of variability. Please see table 2, which summarizes rainfall and growing season temperatures from 1994-1997. The environmental conditions described above could result in larger standard deviations. The variation associated with year to year factors is normally higher than the variation associated within locations or from location to location in a given year.

Table 2. Average temperatures (Fahrenheit) and rainfall (inches) for central Iowa.

TEMPERATURE

YEAR	MAY	JUN	JULY	AUG	AVERAGE
1994	59.8	70.7	71.9	69.0	67.9
1995	56.2	69.4	74.3	76.9	69.2
1996	56.2	69.3	71.3	70.5	66.8
1997	53.5	70.6	74.1	69.6	67.0
AVG	56.4	70.0	72.9	71.5	67.7

RAINFALL

YEAR	MAY	JUN	JULY	AUG	Total
1994	3.67	5.75	1.71	4.18	15.31
1995	5.04	4.19	2.94	2.87	15.04
1996	8.47	4.35	2.51	2.14	17.47
1997	4.32	3.27	4.10	1.36	13.05
AVG	5.38	4.39	2.82	2.64	15.22

ADDITIONAL DESCRIPTION OF PH02T INBRED PER SE YIELD TEST COMPARISON OF PH02T AND PHBG4 EVALUATED OVER YEARS EXHIBIT D.

VARIETY #1 = PH02T VARIETY #2 = PHBG4

1% SIG	R DRP T EAR S ABS	 	5 100.0 1 99.2 4 3 4 6 4 0.8 7* .285	6 98.9 0 100.0 3 2 3 2 4 1.1 3 .500	0 99.6 8 99.5 0 5 2 0 1 2 0 1 3 * .919
# !	BAR PLT ABS		96 4	98. 100. 1.	
SIG	GRN APP ABS	7.5 7.3 7.3 7.0 7.9 7.95	6.7 8.3 3 6 1.6	6.0 7.5 4 4 1.5 1.5	6.6 7.7 9 14 1.2 1.27
ון ייי ייי	BRT STK ABS	97.5	92.2 93.2 55 7 1.0	96.8 97.0 4 5.0 5.0 8999	95.7 97.0 16 21 1.3
SIG +	STK LDG ABS	95.2 76.1 5 10.1 .057*	94.1 77.4 3 6 16.7	94.9 85.7 9.2 9.2	94.8 79.1 11 19 15.7
1 % 1 0% 1 17	STA GRN ABS	6.3 4.8 13 18 1.5	7.0 3.9 11 14 3.1	7.1 4.6 8 2.5	6.7 4.4 32 40 2.3
 	RT LDG ABS	000000000000000000000000000000000000000	90.00 9.00.0 8 4 4.11.236	100.0	94.6 000.0 5 7 7.216
]]] [GDU SLK ABS	1584 1 1564 1 19 19 20 .000#	1582 1553 1 43 47 29 .000#	1607 1 1558 1 30 32 49 .000#	1590 1557 92 98 33
1 1 1 1	GDU SHD ABS	1566 1548 19 19 18	1563 1534 46 51 29	1591 1552 30 32 39	1573 1542 95 102 30
; ; ; ;	EST CNT ABS	40.2 39.5 26 33 0.7	31.8 30.4 32 51 1.4	32.1 31.0 23 30 1.1	34.6 33.5 81 114 1.1
 	SDG VGR ABS	5. 4 4. 8 1. 16 0. 8 4. 8 4. 4 4. 8	6.1 4.3 25 28 1.8	5.9 4.3 14 1.6 .002#	5.9 4.4 55 64 1.5
1 1 1	GQU /HA %MN	128 107 10 20 20 21 21	144 124 124 4 7 20 101	136 113 4 4 23 329	133 112 18 31 21 21
! ! !	GQU /HA ABS	68.4 58.8 10 20 9.6	45.6 38.8 4 7 6.8	35.9 30.8 30.8 4 4 320	56.1 48.1 18 31 8.0
; ! ! !	TST WT ABS	57.5 57.9 10 20 0.4	58.7 58.4 3 6 0.3	53.8 54.1 4 0.3	56.9 57.1 17 30 0.3
; ; ; ;	MST	21.1 21.0 14 24 0.1	18.7 18.4 4 7 0.3	28.0 28.0 44.0 812.0	22.0 21.8 22 35 0.1
i 1 1	BU ACR %MN	128 107 10 20 21 21	144 124 124 4 7 20	136 113 4 4 23 23	133 112 18 31 21 21
! ! !	BU ACR ABS	109.1 93.8 10 20 15.3	72.6 61.9 4 7 10.7	57.3 49.1 4 8.2 320	89.5 76.7 18 31 12.8
 	I VAR #	1 2 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF PR > T	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF
 	YEAR REGION VAR	SUM	SUM	SUM	TOTAL SUM
 	YEAR	94	و ری	96	TOTA

*PR > T values are valid only for comparisons with LOCS >= 10.

D.E.	EDDODUCE LOCALLY Include force purchas and edition data on all source ductions	FORM ARE	DDOVED OND NO 0504 0055
KE	PRODUCE LOCALLY, Include form number and edition date on all reproductions. U.S. DEPARTMENT OF AGRICULTURE	The following statements are made in accord	
	AGRICULTURAL MARKETING SERVICE EXHIBIT E	1974 (5 U. S. C. 552a) and the Paperwork i	•
	STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determinentificate is to be issued (7 U.S.C. 2421). In until certificate is issued (7 U.S.C. 2426).	
1.	NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
	PIONEER HI-BRED INTERNATIONAL, INC.	OR EXPERIMENTAL NUMBER	РН02Т
4	.ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
	7301 NW 62 nd AVENUE	515-270-4051	515-253-2125
	P.O.BOX 85	7. PVPO NUMBER	-
	JOHNSTON, IA 50131-0085		
8.	Does the applicant own all rights to the variety? Mark an "X" in appropriate bl	ock. If no, please explain:	YES NO
		-	
_	to the applicant (individual or company) a LLC mational and C. become		VES EL NO
9.	Is the applicant (individual or company) a U.S. national or U.S. based compan	ıy? ⊠	YES NO
10.	If no, give name of country Is the applicant the original owner? ✓ YES ☐ NO	If no, please answer one of the followin	ng:
	a. If original rights to variety were owned by individual(s), is (are) the origin	-	-
	☐ YES ☐ NO	if no, give name of country	
	b. If original rights to variety were owned by a company(ies), is(are) the ori		
	⊠ YES □ NO	If no, give name of country	
11.	Additional explanation on ownership (if needed, use reverse for extra space):		
	PH02T is owned by Pioneer Hi-Bred International, Inc.		
TOT T	ZAON NORTH		
	EASE NOTE:		
Plar	nt variety protection can be afforded only to owners (not licensees) who meet one of the	e following criteria:	
1.	If the rights to the variety are owned by the original breeder, that person must be a U Which affords similar protection to nationals of the U.S. for the same genus and specific		ntry, or national of a country
2.	If the rights to the variety are owned by the company which employed the original brocountry, or owned by national of a country which affords similar protection to national		
3.	If the applicant is an owner who is not the original owner, both the original owner and	d the applicant must meet one of the above crit	eria.
The	original breeder/owner may be the individual or company who directed final breeding.	See section 41(a)(2) of the Plant Variety Prot	ection Act for definition.
Δ	ccording to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of info	rmation unless it displays a valid OMB control number. The	ne valid OMB control number for this

According to the Paperwork Reduction Act or 1995, no persons are required to a collection of information unless it displays a valid OMB control number. The valid OMB control number is of the information in 1998 of the information is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD) USDA is an equal employment opportunity employer.